

Dissipative maps at the chaos threshold: Numerical results for the single-site map

Ugur Tirnakli¹

Department of Physics, Faculty of Science, Ege University, 35100 Izmir, Turkey

Abstract

We numerically study, at the edge of chaos, the behaviour of the single-site map $x_{t+1} = x_t - x_t/(x_t^2 + \gamma^2)$, where γ is the map parameter.

Key words: Nonextensive thermostatics, dynamical systems

PACS: : 05.45.-a, 05.20.-y, 05.70.Ce

In recent years, a number of studies has addressed the nonlinear dynamical systems at the chaos threshold [1–8]. At this special point (and also at some others such as period doublings, tangent bifurcations points where the Lyapunov exponent vanishes), it is shown that the proper entropy appears to be the nonextensive form [9] $S_q = k \left(1 - \sum_{i=1}^W p_i^q\right) / (q - 1)$ with a proper q value (say q^*) for each map under consideration. Up to now, three different methods are available for the determination of the q^* values. First method comes from the analysis of the sensitivity to initial conditions where a general sensitivity function

$$\xi(t) = [1 + (1 - q)\lambda_q t]^{1/(1-q)} \quad (q \in R), \quad (1)$$

can be introduced. Here, λ_q is the generalized Lyapunov exponent and this form recovers the standard exponential one for $q = 1$, but it generically exhibits a power-law behaviour. In fact, eq.(1) corresponds to the power-law growth of the upper bound of the sensitivity function and this upper bound allows us to determine the proper q^* value of the studied map. Second method is based on the geometrical aspects of the attractor at the edge of chaos and uses the scaling behaviour of the multifractal singularity spectrum $f(\alpha)$ as

$$\frac{1}{1 - q^*} = \frac{1}{\alpha_{min}} - \frac{1}{\alpha_{max}} \quad (q^* < 1) \quad (2)$$

¹ tirnakli@sci.ege.edu.tr

where α_{min} and α_{max} being the most concentrated and most rarefied regions of the measure on the attractor respectively. It is obvious that calculating these values from the $f(\alpha)$ curve, one can easily estimate the proper q^* value [3]. Finally, third method, which was introduced very recently, deals with the entropy increase rates [5,6]. It basically based upon the (ensemble version) of generalized Kolmogorov-Sinai entropy $K_q \equiv \lim_{t \rightarrow \infty} \lim_{W \rightarrow \infty} \lim_{N \rightarrow \infty} [S_q(t)]/t$ and it is conjectured that (i) a special q^* value exists such that K_q is finite for $q = q^*$, vanishes for $q > q^*$ and diverges for $q < q^*$; (ii) this q^* value coincides with that coming from the other two methods explained above.

Now we are prepared to introduce the main purpose of the present work. Here, we shall try to verify once again these conjectures using the abovementioned three methods for the single-site map

$$x_{t+1} = x_t - \frac{x_t}{x_t^2 + \gamma^2} \quad , \quad (3)$$

where γ is the map parameter [10]. The shape of the map is illustrated in Fig. 1. This map has a route to chaos by period doubling with Feigenbaum scaling but it exhibits an interesting property that the bifurcation cascade splits into two independent ones, which both obey Feigenbaum scenario. This property is the generic peculiarity for the single-site map in contrast to the logistic map [10]. On the other hand, these two independent cascades obey Feigenbaum scaling, which means that the Feigenbaum numbers of this map are the same as those of the logistic map, therefore, in that sense the single-site map shares the same universality class with the logistic map. Consequently, what we expect is that, at the chaos threshold, the proper q^* value of this map would coincide with the value of the logistic case (namely, $q^* \simeq 0.24$). Our numerical results check and verify this expectation. Firstly, let us perform the first method to estimate the proper q^* value of the single-site map at the onset of chaos (namely at $\gamma_c = 0.43400483\dots$). For the extremal point of the map ($x_0 = 0.3213\dots$), the sensitivity function exhibits a power-law divergence, $\xi(t) \propto t^{1/(1-q)}$ as shown in Fig. 2. Measuring the upper bound slope $1/(1-q)$, we determine the proper q value as $q^* = 0.24$. Now, let us proceed with the second method. In Fig. 3, the $f(\alpha)$ curve is given, where the monotonic behaviour with respect to the number of iterations is evident. Since the correct values of α_{min} and α_{max} could be obtained in principle when the number of iterations goes to infinity, the extrapolation shown in the inset of Fig. 3 is used to estimate the correct values of α_{min} and α_{max} . Using these values in the scaling relation Eq.(2), it is found that the proper q value is again $q^* = 0.24$. (We also verify using the extrapolation technique that the fractal dimension d_f of this map coincides with that of the logistic map as expected). Finally, we shall use the third method for this map. To do this, we partition the phase space, $x \in (-1/2\gamma, 1/2\gamma)$, into W equal cells, then we choose one of them and select N initial conditions all inside the chosen cell. As time

evolves, these N points spread within the phase space and finally this yields a set of probabilities. In the beginning of time, of course, $S_q(0) = 0$, then it exhibits three successive regions. In the intermediate region, the entropy starts increasing and this is the region where the linear increase of the proper entropy is expected to emerge (for details see [5,6]). In Fig. 4, we plotted the time evolution of the entropy for three values of q . (At this point, it should be mentioned that the procedure of averaging over the efficient cells as described in [5,6] has also been performed here). In the intermediate region, it is seen that the linear increase of the entropy with time emerges for a special value of q . To support this picture, we fit the curve with the polynomial $S_q = A + Bt + Ct^2$ in the intermediate region $[t_1, t_2]$. Since the nonlinearity coefficient $R \equiv C(t_1 + t_2)/B$ is a measure of the importance of the nonlinear term, it should vanish for a strickly linear fit. Therefore, inset of Fig. 4, we estimate the proper q value as $q^* = 0.234$, which is the value when R is zero.

Summing up, we study the behaviour of the single-site map at the chaos threshold. We numerically verify that the proper entropic index for this map is $q^* \simeq 0.24$. This constitutes another example of the one-dimensional dissipative maps which behave as $q \neq 1$ at the chaos threshold.

This work has been supported by Ege University Research Fund under the project number 2000 FEN 049.

References

- [1] C. Tsallis, A.R. Plastino and W.-M. Zheng, Chaos, Solitons and Fractals **8** (1997) 885.
- [2] U.M.S. Costa, M.L. Lyra, A.R. Plastino and C. Tsallis, Phys. Rev. E **56** (1997) 245.
- [3] M.L. Lyra and C. Tsallis, Phys. Rev. Lett. **80** (1998) 53.
- [4] U. Tirnakli, C. Tsallis and M.L. Lyra, Eur. Phys. J. B **11** (1999) 309.
- [5] V. Latora, M. Baranger, A. Rapisarda and C. Tsallis, Phys. Lett. A **273** (2000) 97.
- [6] U. Tirnakli, G.F.J. Ananos and C. Tsallis, Phys. Lett. A (2001), in press. (cond-mat/0005210).
- [7] U. Tirnakli, Phys. Rev. E **62** (2000) 7857.
- [8] F.A.B.F. de Moura, U. Tirnakli and M.L. Lyra, Phys. Rev. E **62** (2000) 6361.
- [9] C. Tsallis, J. Stat. Phys. **52** (1988) 479.
- [10] S.L. Ginzburg and V.I. Sbitnev, Physica D **132** (1999) 87.

Figure Captions

Figure 1 - The shape of the single-site map with $\gamma = 0.3$.

Figure 2 - Log-log plot of the sensitivity function versus time.

Figure 3 - Approximate multifractal singularity spectrum of the critical attractor of the single-site map for increasingly large number of iterations, going from $N = 64$ to 4096. Inset: α_{min} , d_f and α_{max} from the singularity spectra obtained from distinct number of iterations. Extrapolated values of them are also given.

Figure 4 - Time evolution of $S_q(t)$ for three different values of q . $W = 10^5$ and $N = 10^6$ is used. The interval characterizing the intermediate region is $[9,21]$. Inset: The nonlinearity coefficient R versus q (see text for details).

